

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Appendix F AMENDED DRAWINGS




FIGURE 1




FIGURE 2




FIGURE 13







FIGURE 14




or



$R' =$




or



where * is the point of attachment.

FIGURE 16



induced excited state
 ✓
 Optical limiting by two-photon absorption in 4-dimethylamino-4'-dibutylaminostilbene (MBDAS) (points). Straight line corresponds to the linear transmission of 96%.

FIGURE 17





FIGURE 19

Linear transmission spectrum (solid line) of a mixture of 4,4'-bis(dibutylamino)stilbene and 4,4'-bis(diphenylamino)stilbene for a total molecular concentration of 0.1M. Spectra of effective two-photon absorption cross sections for 4,4'-bis(di-n-butylamino)stilbene (triangles) and 4,4'-bis(diphenylamino)stilbene (squares).



Nonlinear transmission data showing enhanced two-photon absorption in a
 1-(4-dimethylaminophenyl)-10-(4-dibutylaminophenyl)-deca-1,3,5,7,9-pentane
 (squares) as compared to the ^{effective} ~~substituted BODAP~~ ^{4,4'-bis(di-n-butylamino)stilbene} (circles). Measurements were
 performed at 598 nm using 0.01M solutions.

FIGURE 20